

Potential for woody plant control by Spanish goats in the sagebrush steppe

B. Fajemisin^a, D. Ganskopp^{b,*}, R. Cruz^c, M. Vavra^d

^a National Animal Production Research Institute, Ahmadu Bello University, Zaria, Nigeria

^b USDA Agricultural Research Service, HC-71 4.51 Hwy. 205, Burns, OR 97720, USA

^c Department of Rangeland Resources, Oregon State University, Corvallis, OR 97731, USA

^d Eastern Oregon Agricultural Research Center, HC-71 4.51 Hwy. 205, Burns, OR 97720, USA

Accepted 7 August 1995

Abstract

Botanical composition of the diet of eight Spanish goats was studied on the Northern Great Basin Experimental Range during two stages of plant phenology (active growth in early July and cured forage in mid-August) to assess their potential for the control of sagebrush (*Artemisia tridentata* ssp. *wyomingensis* Nutt.) and western juniper (*Juniperus occidentalis* Hook.) and their nutritional status on sagebrush-steppe rangelands. Diets were quantified by documenting the number of visits to each species, the bites harvested and time expended grazing each forage over 4 consecutive days in each period. Forage chemical characteristics evaluated included: CP, NDF, ADF, ADL and IVDMD. Diet composition and forage quality indices varied significantly ($P < 0.05$) with changes in plant phenology. Vegetation cover averaged 52% in the pasture and consisted of 36% grasses, 8% forbs, 7% shrubs and 0.6% trees. Available herbage (excluding woody plants) was 534 kg ha⁻¹ during active growth trials and 572 kg ha⁻¹ when forages had cured. When forages were green goats acquired 28% of their total bites from grasses, 71% from forbs, 0.3% from shrubs and 0.9% from juniper trees. After forages had cured values were 35% from grasses, 56% from forbs, 0.1% from shrubs and 8.8% from juniper. The browsing of juniper (both foliage and bark), after herbaceous forages had cured, was the only substantive use of woody plants. Sagebrush was only lightly used (0.2% of total bites) when herbaceous forages were actively growing. Available herbage was of relatively high quality during both trials. When forages were actively growing, CP of grasses ranged from 8.9 to 5.6%, forbs from 17.7 to 8.5%, sagebrush scored at 8.5%, juniper foliage at 8.1% and juniper bark at 3.2%. After herbaceous forages had cured, grass CPs ranged between 6.2 and 3.1%, forbs from 10.4 to 4.4%, shrubs were not sampled and juniper foliage averaged 7.6%. Given the low levels of browsing exhibited by goats on sagebrush and juniper we see little opportunity for control of these woody plants when pastures provide a diverse ($N = 25$ species) array of readily available (534–572 kg ha⁻¹) and nutritious forages and pastures are lightly stocked (0.63 goats ha⁻¹ month⁻¹). We do need, however, to further explore their potential for control of these species under several other regimes. These include extended trials on more deteriorated rangelands, trials during seasons or conditions where forage is limited and the animal's selective opportunity is restricted and trials in high quality environments on newly established sagebrush and juniper seedlings.

Resumen

La composición botánica de las dietas de 8 cabras españolas fue evaluada en el Northern Great Basin Experimental Range durante 2 etapas de fenología vegetal (crecimiento activo a principios de Julio y dormancia a mediados de Agosto)

* Corresponding author. Tel.: 503-573-2064; fax: 503-573-3042; e-mail: ganskopp@bcc.orst.edu.

para evaluar su potencial en el control de sagebrush (*Artemisia tridentata* sspp. *wyomingensis* Nutt.) y western juniper (*Juniperus occidentalis* Hook.), así como el estado nutricional de estas en praderas en el sagebrush steppe. Las dietas fueron cuantificadas observando el número de visitas, el número de mordidas, y el tiempo que duraron pastoreando cada forraje durante cuatro días consecutivos. Las características químicas de los forrajes evaluadas incluyeron: CP, NDF, ADF, ADL, y IVDMD. La composición de la dieta y la calidad de los forrajes variaron significativamente ($P < 0.05$) con los consistiendo de 36% de pastos, 8% de hierbas, 7% de arbustos y 0.6% de árboles. El forraje disponible (excluyendo plantas leñosas) fue 534 kg ha^{-1} durante las pruebas en la etapa de crecimiento activo y 572 kg ha^{-1} en la etapa de dormancia. Cuando los forrajes estaban verdes las cabras obtuvieron del total de las mordidas 28% de pastos, 71% de hierbas, 0.3% arbustos, y 0.9% de juníperos. El pastoreo de junípero (tanto follaje como corteza) después de que los forrajes estaban en dormancia, fue el único uso substancial de plantas leñosas. El sagebrush solo fue consumido ligeramente (0.2% de las mordidas totales) cuando las plantas herbáceas estaban en crecimiento activo. El forraje disponible fue de calidad relativamente alta durante las dos etapas. Cuando los forrajes estaban en crecimiento la CP de los pastos osciló entre 8.9–5.6%, las hierbas entre 17.7–8.5%, sagebrush 8.5%, follaje de junípero 8.1% y corteza de junípero 3.2%. Después que los forrajes habían madurado, la CP de los pastos fue entre 6.2–3.1%, hierbas entre 10.4–4.4%, los arbustos no fueron muestreados, y el follaje de junípero promedio 7.6%. Dados los bajos niveles de consumo de sagebrush y junípero exhibidos por las cabras vemos pocas oportunidades para el control de estas plantas leñosas cuando las praderas proveen una combinación diversa ($N = 25$ especies) de forrajes nutritivos disponibles ($534\text{--}572 \text{ kg ha}^{-1}$) en praderas con baja carga animal (0.63 cabras/ha/mes). Sin embargo, debemos explorar el potencial de las cabras para el control de estas especies bajo diferentes circunstancias. Estas incluyen pruebas prolongadas en praderas más deterioradas, pruebas durante períodos cuando el forraje es limitado y las oportunidades de selección de los animales es restringida, y praderas de alta calidad con juníperos y sagebrush recién establecidos.

Keywords: Biological control; Sagebrush; Juniper; Forage selection; Grazing behavior; Nutrition; Forage quality

1. Introduction

Owing to a variety of factors, the Northern Great Basin sagebrush steppe is experiencing expansion of western juniper (*Juniperus occidentalis* Hook.) stands and increases in the density and cover of sagebrush (*Artemisia* spp.) on otherwise good condition rangelands (Miller et al., 1994). In both instances this is occurring at the expense of perennial grasses and forbs. Chemical controls are becoming prohibitively expensive and politically unpopular (Britton and Sneva, 1979; Bastian et al., 1995), and biological efforts have shown little potential to date for control of these competitive plants (Richman et al., 1994). While severe defoliation can kill big sagebrush (Pringle, 1960; Cook and Stoddart, 1960; Wright, 1970) the literature provides no information on seasons or intensities of defoliation required to affect western juniper. This being the case, many avenues for biological control of these woody plants remain largely unexplored, including use of the various breeds of goats.

Use of livestock in prescription grazing programs to control undesirable plants demands knowledge of the animal's innate dietary preferences and the com-

position and nutritive value of the forage resources. Both aspects must be integrated in biological control efforts to affect the target plant without impacting desirable components of the vegetation and to assure adequate livestock production. Among ruminants, the goat (*Capra hircus*) is well known for utilizing a broad array of forages which explains their ability to adapt to various environments, especially the arid areas (Morand-Fehr, 1988). Leclerc (1984) reported that browse constituted 75% to the diet of goats in the Corsican 'maquis' during winter but did not exceed 15% during spring. Similarly, Narjisse and El Bare (1986) found woody plants contributed most significantly to the diet (68%) when herbaceous forage was less available on the forest floor. Goats appear to be especially flexible in their food selection and can maximize nutritional benefits through their selective grazing habits. The objectives of this study were (1) to assess the potential of Spanish goats for woody plant control during both growing and dormant forage conditions on good condition sagebrush rangeland, and (2) to evaluate the nutritive value of the goats dietary choices on northern Great Basin, sagebrush steppe vegetation. Direct observation techniques were used to quantify the relative

amounts of plant species selected by goats as they freely grazed during two 4-day trials. One trial was conducted in early July, when all forages were actively growing and of high nutritional value. The second was conducted in mid-August when herbaceous vegetation had ceased growing and cured to marginal nutritional levels.

2. Materials and methods

The study was conducted in 1993 on the Northern Great Basin Experimental Range about 65 km southwest of Burns, Oregon. This semi-arid desert range (elevation 1370 m) lies in the northern edge of the Great Basin and receives a mean annual precipitation of about 30 cm. Approximately 80% of the moisture falls between October and June with the majority occurring as snow. Growth of herbaceous forages is typically arrested by late July, as soil moisture is gradually depleted by evapotranspiration during the growing season.

Tree, shrub, and herbaceous plant density and cover were sampled in a temporary, electrically fenced, 1.7 ha pasture prior to the July grazing trial. The pasture has a long history (50+ years) of brief summer use by cattle, and with the exception of increasing sagebrush and juniper, supported a very desirable and productive array of native grasses and forbs. Foliage cover of each species was measured along five 50-m line transects systematically arranged within the pasture. Frequency of occurrence and density of shrubs and herbaceous plants were sampled from 40 1-m² randomly distributed plots, and all western juniper trees in the pasture were counted. Available biomass of herbaceous plants was harvested from 25 1-m² plots prior to each of the two grazing trials and oven dried at 60°C. Grasses were sorted by species in each plot while forbs were composited. Herbage production of trees and shrubs was not sampled. Prior to each trial additional samples of the most prominently utilized forages were harvested, oven dried, ground to pass a 1-mm screen, and retained for later chemical analyses of forage quality.

Eight dry, non-pregnant, adult female goats, weighing 26.3 ± 1.2 kg (mean \pm SE), were grazed to quantify plant species selected over 4 consecutive

days during each of two forage phases: actively growing (July 6–9) and dry, dormant forages (August 17–20). Although the animals were on station approximately 6 weeks before the trials, we elected to limit their exposure to native rangeland to the 24 h period immediately prior to sampling when they were confined in a temporary holding paddock in one corner of the pasture. The paddock supported sagebrush, juniper, several of the grasses common to the pasture, and a portion of the forb component. We followed this protocol because the goat industry is not well established in our region, and local land managers seeking to implement biological controls would typically be forced to import herds of naive animals from other areas.

Goats were released in a random sequence in the morning and dietary composition estimated by direct observation and tally of individual bites (Altman, 1974; Holechek et al., 1984; Austin et al., 1994; Jones et al., 1994; Richman et al., 1994). A technician equipped with a laptop computer recorded each bite harvested by pressing the space bar, and noted walking or transition between feeding stations by pressing the 'W' key. The subsequent data file consisted of a series of time entries (i.e. 09:32:17.6), each denoting an individual bite, and interjections where the 'W' key was pressed (i.e. W09:32:18.1 126) which referenced the time of movement to a new feeding station and the observation number (126) of the next plant to be grazed. A second technician concurrently recorded the species of each plant as it was grazed. Each goat was monitored as it foraged upon a series of 50 plants. Sampling typically occupied the entire morning, and the goats were allowed to remain in the pasture until evening when they were penned overnight to avoid potential coyote (*Canis latrans*) predation.

Integration of bite-count files and the concurrently generated species lists allowed derivation of several variables for each animal. These included: the number of visits to each species, the total number of bites harvested from each, the mean number of bites per visit to each forage, and calculation of the mean time expended grazing each species (derived from the times of the first and last bites at each feeding station). Software for acquisition and integration of these data were generated in GWBasic by the second author.

Analyses of forage quality samples included: dry matter, organic matter, ash content, and Kjeldahl nitrogen as determined by procedures of the Association of Official Analytical Chemists (1980). NDF, ADF, and ADL were measured according to Goering and Van Soest (1970), and IVDMD was determined using techniques of Tilley and Terry (1963).

With the exception of 'visits to plants', which is autocorrelated because the same number of observations occurred for each goat or block, variables describing grazing behavior of goats were analyzed using a split-split plot analyses of variance. Data indexing total bites harvested and total time expended grazing each species were converted to percentages for presentation after analyses. Individual goats ($N = 8$) functioned as replicates or blocks. Stages of phenology ($N = 2$) and days ($N = 4$) could not be randomized and served as whole plots and subplots, respectively, and plant species ($N = 25$) functioned as sub-sub plots. Mean separations were accomplished with Fisher's protected LSD procedures ($P < 0.05$). Total number of observations in the two trials was 1600 (8 animals \times 2 stages of phenology \times 4 days \times 25 forage species).

3. Results and discussion

Herbaceous vegetation covered roughly 45% of the soil surface. Idaho fescue (*Festuca idahoensis* Elmer) ranked highest in cover and occurred with the greatest frequency, being present in 95% of the plots sampled (Table 1). Sandberg's bluegrass (*Poa sandbergii* Vasey), however, attained the highest density with roughly 4.5 plants m^{-2} . Among the forbs, bushy birdbeak (*Cordylanthus ramosus* Nutt.) ranked highest in cover (3.3%), density (3.4 m^{-2}), and frequency (80%). The shrub layer was dominated by Wyoming big sagebrush (*Artemisia tridentata* ssp. *wyomingensis* Beetle), and western juniper characterized the overstory with a density of 169 trees ha^{-1} . Approximately 44% of the junipers were less than 2 m tall with the remaining being scored as mature trees. Because ground level branches are persistent on these trees, foliage of both juvenile and mature specimens is within easy reach of browsing goats.

Precipitation during the preceding September–

Table 1

Characteristics of the sagebrush steppe vegetation grazed by Spanish goats on the Northern Great Basin Experimental Range in 1993. Nomenclature according to Hitchcock and Cronquist (1973)

Species	% Cover	Density (m^{-2})	% Frequency
<i>Festuca idahoensis</i>	15.7(1.9)	3.8(0.4)	95
<i>Agropyron spicatum</i>	7.6(1.5)	1.4(0.2)	63
<i>Poa sandbergii</i>	7.3(1.2)	4.5(0.5)	83
<i>Koeleria cristata</i>	4.2(1.8)	1.8(0.3)	70
<i>Stipa thurberiana</i>	0.9(1.8)	0.6(0.3)	20
<i>Sitanion hystrix</i>	0.3(0.1)	0.2(0.1)	13
<i>Bromus tectorum</i>	0.1(0.0)	0.2(0.1)	13
Total grasses	36.1	12.5	
<i>Cordylanthus ramosus</i>	3.3(0.9)	3.4(0.5)	80
<i>Eriogonum spheroccephalum</i>	1.6(1.2)	0.1(0.1)	10
<i>Erigeron linearis</i>	1.1(0.7)	1.0(0.2)	45
<i>Erigeron filifolius</i>	0.6(0.3)	0.3(0.1)	23
<i>Phlox hoodii</i>	0.5(0.2)	0.1(0.1)	28
<i>Astragalus filipes</i>	0.5(0.2)	0.5(0.1)	10
<i>Lupinus caudatus</i>	0.4(0.2)	–	35
<i>Achillea millefolium</i>	0.3(0.3)	0.2(0.1)	5
<i>Crepis acuminata</i>	0.1(0.0)	–	10
<i>Penstemon speciosus</i>	0.1(0.1)	0.2(0.2)	5
Total forbs	8.4	5.8	
<i>Artemisia tridentata</i>	6.2(0.3)	3.0(0.1)	
<i>Chrysothamnus viscidiflorus</i>	0.4(0.2)	–	
Total shrubs	6.5	3.0	
<i>Juniperus occidentalis</i>	0.6(0.4)	169.3/ha	
Total trees	0.6	169.3	

–, not encountered during sampling, or only negligible quantities found. Values are means, with standard errors given in parentheses.

June period is correlated strongly with annual yield of herbaceous forage in the region (Sneva, 1982) and was 167% of the long-term mean (255 mm) for the 1993 forage year. As a result, forage availability was greater than average. Among the grasses, Idaho fescue and bluebunch wheatgrass (*Agropyron spicatum* Scribn. & Smith) ranked highest in herbage production (Table 2). Availability of grasses decreased approximately 14% between the active growth and cured forage sampling periods, while the later maturing forbs increased in availability from 41.2 to 151.2 $kg\ ha^{-1}$ in the same period. The increase in forb biomass more than offset the decrease in grasses, and

Table 2

Available herbage of sagebrush steppe vegetation grazed by Spanish goats on the Northern Great Basin Experimental Range during active growth and cured forage stages of phenology in 1993

Species	Active growth— July(kg ha ⁻¹)	Cured forage— August(kg ha ⁻¹)
<i>Festuca idahoensis</i>	131.6(23.2)	122.8(20.1)
<i>Agropyron spicatum</i>	111.6(30.2)	141.6(38.5)
<i>Poa sandbergii</i>	123.8(21.5)	58.8(8.1)
<i>Koeleria cristata</i>	67.8(7.6)	66.4(13.3)
<i>Stipa thurberiana</i>	22.6(11.8)	21.6(9.0)
<i>Sitanion hystrix</i>	19.6(10.4)	6.0(3.5)
<i>Bromus tectorum</i>	16.0(6.7)	4.4(2.1)
Total grasses	493.0	421.6
Total forbs	41.2(14.7)	151.2(31.5)

Values are means with standard errors given in parentheses.

implied that total available herbaceous material increased from 534 to 572 kg ha⁻¹ between the green and dormant stages of growth.

Although we did not sample shrub and tree biomass, Richman (1993) quantified sagebrush production in a similar environment in 1990 and 1991. Richman (1993) estimated sagebrush cover at 4.3% and twig and leaf biomass at 340 kg ha⁻¹. Given our measure of 6.2% sagebrush cover and more precipitation than Richman's 1990–1991 sampling period, we estimate our sagebrush biomass at least equaled or exceeded 486 kg ha⁻¹. We have no means of estimating availability of juniper foliage.

In our analyses of the goats foraging activities, forage species and species × phenology effects were consistently significant ($P < 0.01$) for all variables, while days and interactions with days were not. Consequently, data are presented for each stage of phenology (Table 3). Forbs were the most prominent dietary constituent during both stages of phenology. Grasses ranked second, while shrubs and trees were of minor importance.

When forages were actively growing, forbs and grasses constituted 64.0% and 33.7%, respectively, of the total plants visited (Table 3). The mean number of bites monitored per goat in a daily grazing session was 169 with roughly 64% harvested from forbs and 34% selected from grasses. Shrubs and trees were minor dietary components. Mean foraging time measured for each goat averaged 5.7 min day⁻¹

with approximately 69% and 29% devoted to forbs and grasses, respectively. Since time involved in travel and scrutiny of forages was not included in this measure, actual periods of observation were considerably greater in duration.

During the dormant stage of phenology, visits to forbs still dominated (50.6%), but they were less prominent in the diet than when forages were green. Visits to grasses increased roughly 12% over the earlier sampling period, and the number of bites harvested from grasses increased by slightly over 6%. On average, each goat was monitored for 200 bites and roughly 5.3 min of actual foraging time per day. Again, shrubs were of minor importance, but bites harvested from juniper trees did increase to nearly 9% of the total.

These observations mirror those of several others. Fedele et al. (1988) and Pizzillo et al. (1988) found the spring diets of goats were primarily grasses (65–75%), followed by other non-grass and non-legume families (22–25%) and legume families (3–10%). Grasses diminished in the diet during summer (55–60%), forbs increased (40–45%) and the legume component tended to disappear (Fedele et al., 1988; Pizzillo et al., 1988).

Shrubs appeared to be the class of plants least preferred by goats during both trials, as indicated by low values for visits to plants (0.8–0.2%) and percentage of total bites (0.1–0.3%). Although not presented in our data summaries, the highest number of bites per visit (8.6–10.8) and time per visit (16.0–29.8 s) were recorded for *Elymus cinereus*. This was a large-stature grass (over 2 m high), and was the only graminoid from which the goats foraged by stripping individual leaves from the plant. *Astragalus filipes* and *Achillea millefolium* were the most visited forbs. Among the forbs the greatest number of bites per visit (5.3–6.2) occurred with *Lithospermum ruderae* which was quite rare and remained succulent throughout both trials. While the patterns of visits to and bite counts on individual plants did not generally change much between trials, there was considerable variation among the forage classes.

Chemical composition and in-vitro organic matter disappearance for plants grazed by goats varied considerably between the green and dormant stages of phenology. Levels of various plant nutrients, constituting 5% or more of the total bites at either stage of

growth, are listed in Table 4. Collectively, the forages in Table 4 accounted for roughly 86% and 91% of the total bites harvested during the active growth and dormant stages of phenology, respectively. Crude protein and IVDMD values were consistently higher for forages in the first period than in the second, and generally, the reverse occurred for DM, NDF, ADF, ADL and ash values. Weighting of CP values of forages by the proportion of total bites taken from

each (Schwartz et al., 1977) yielded CP estimates of the goats diet of 10.7% during active growth and 6.7% when forages were cured. These values are most certainly negatively biased, however, because grazers typically harvest diets of higher quality than our best hand compounded efforts.

Studies of seasonal composition of goat diets by Malechek and Provenza (1981) and Rubino et al. (1988) have shown that the ration, especially on arid

Table 3

Percentage of total visits, total bites harvested, and total grazing time expended on each forage species by Spanish goats grazing sagebrush steppe vegetation for 4 days periods during two stages of phenology on the Northern Great Basin Experimental Range in 1993

Species	Active growth—July			Cured forage—August		
	% Visits	% Bites	% Time	% Visits	% Bites	% Time
<i>Agropyron spicatum</i>	12.3	8.9d	8.8abcd	17.3	11.6ef	10.3cde
<i>Agropyron desertorum</i>	7.7	8.1d	9.0bcd	8.8	7.9cde	8.2cd
<i>Festuca idahoensis</i>	7.9	5.0bcd	4.1abc	7.2	4.2bc	7.7c
<i>Elymus cinereus</i>	0.8	2.6ab	3.9abc	0.9	2.2ab	2.2ab
<i>Poa sandbergii</i>	3.1	2.5ab	2.4abc	0.9	0.6ab	0.4a
<i>Sitanion hystrix</i>	0.4	0.2a	0.2a	1.2	0.5ab	0.4a
<i>Koeleria cristata</i>	0.6	0.5a	0.4ab	8.7	7.6cd	5.9bc
<i>Bromus tectorum</i>	0.8	0.4a	0.4ab	0.0	0.0a	0.0a
<i>Stipa thurberiana</i>	0.1	0.0a	0.0a	0.3	0.1a	0.2a
Total grasses	33.7	28.2	29.2	45.3	34.7	35.3
<i>Astragalus filipes</i>	14.4	22.1f	21.8e	19.1	24.7g	23.3f
<i>Achillea millefolium</i>	15.6	16.8e	14.5de	12.9	11.8f	9.1cde
<i>Lepidium perfoliatum</i>	11.6	8.8d	8.5abcd	1.8	1.3ab	1.3a
<i>Crepis accuminata</i>	6.6	6.6cd	6.5abcd	1.7	1.3ab	1.0a
<i>Erigeron filifolius</i>	3.2	2.3ab	2.3abc	1.9	1.5ab	1.2a
<i>Eriogonum spheroccephalum</i>	4.6	3.4abc	2.9abc	1.1	0.6ab	0.6a
<i>Sisymbrium altissimum</i>	1.6	1.4ab	1.5abc	0.3	0.1a	0.1a
<i>Lithospermum ruderae</i>	5.2	8.2d	9.8cd	6.6	11.9f	12.2de
<i>Lupinus caudatus</i>	0.4	0.3a	0.5ab	0.4	0.2a	0.1a
<i>Penstemon speciosus</i>	0.4	0.5a	0.6ab	3.8	2.3ab	1.9ab
<i>Phlox hoodii</i>	0.4	0.4a	0.3ab	0.1	0.1a	0.1a
<i>Cordylanthus ramosus</i>	0.0	0.0a	0.0a	0.9	0.6ab	0.7a
Total forbs	64.0	70.8	69.2	50.6	56.4	51.7
<i>Artemisia tridentata</i>	0.6	0.2a	0.2a	0.1	0.0a	0.0a
<i>Chrysothamnus viscidiflorus</i>	0.1	0.1a	0.4ab	0.1	0.1a 0.3a	
<i>Tetradymia canescens</i>	0.1	0.0a	0.0a	0.0	0.0a	0.0a
Total shrubs	0.8	0.3	0.6	0.2	0.1	0.3
<i>Juniperus occidentalis</i>	1.5	0.9a	1.0ab	3.9	8.8def	12.6e
Total trees	1.5	0.9	1.0	3.9	8.8	12.6

Means in columns sharing a common letter do not differ significantly ($P > 0.05$).

rangelands, undergoes considerable modification depending on climate and year. Among the grasses, *Agropyron desertorum* had the highest crude protein content (8.9–6.2%) and *Agropyron spicatum* the lowest (5.6–4.4%). NDF values for grasses ranged from 58.1 to 71.1%, ADF values from 40.9 to 53.2%, and ADL values from 5.3 to 4.45%. Among forbs, *Astragalus filipes* had highest CP value of 17.7–

7.3%, and *Crepis accuminata* the lowest at 8.5–4.4%. Among the forbs, NDF values ranged from 25.3 to 67.4%, ADF values from 26.8 to 57.0% and ADL values from 4.3 to 10.1%. IVDMD ranged from 44.6 to 63.4% for grasses and from 44.1 to 81.8% for forbs.

The only substantial grazing that occurred on woody plants involved juniper during the dormant

Table 4

Percent DM, CP, NDF, ADF, ADL, IVDMD, and ash content of forages constituting more than 5% of the diet of Spanish goats, as determined from bite count data on the Northern Great Basin Experimental Range during two stages of phenology in 1993. Upper and lower positions of numerical pairs depict data during active growth and cured forage stages of phenology, respectively

Forage class and species	Forage constituent						
	%DM	%CP	%NDF	%ADF	%ADL	%IVDMD	%Ash
Grasses							
<i>Festuca idahoensis</i>	53.3	6.1	58.9	44.3	5.3	60.3	17.1
	61.1	3.1	71.1	52.4	5.8	44.6	19.3
<i>Agropyron spicatum</i>	55.3	5.6	62.1	42.5	6.5	55.4	8.8
	59.6	4.4	63.9	48.8	6.3	48.1	14.3
<i>Koeleria cristata</i>	–	–	–	–	–	–	–
	56.8	5.9	58.1	53.2	5.3	63.4	18.8
<i>Agropyron desertorum</i>	44.8	8.9	66.9	40.9	8.4	56.8	5.9
	61.7	6.2	69.2	43.8	7.2	50.8	8.3
Forbs							
<i>Astragalus filipes</i>	33.4	17.7	52.5	42.4	8.5	54.8	4.8
	45.1	7.3	60.1	47.7	10.1	50.7	5.3
<i>Achillea millefolium</i>	35.1	8.6	33.0	23.1	4.8	80.5	12.6
	51.3	10.4	49.2	36.1	7.1	74.6	16.9
<i>Crepis accuminata</i>	28.1	8.5	30.9	26.8	3.9	81.8	8.2
	45.0	4.4	42.9	36.4	7.7	65.1	9.9
<i>Lepidium perfoliatum</i>	53.0	9.9	51.8	41.1	8.5	54.8	5.2
	86.5	3.9	67.1	57.0	8.8	44.1	7.2
<i>Lithospermum ruderae</i>	31.6	10.5	30.2	25.3	4.3	73.5	23.8
	38.1	7.8	34.6	27.6	4.7	67.5	28.4
Shrubs							
<i>Artemisia tridentata</i>	45.0	8.5	40.3	30.1	6.2	61.3	4.4
	–	–	–	–	–	–	–
Trees							
<i>Juniperus occidentalis</i> (foliage)	58.7	8.1	40.5	29.3	14.2	49.6	3.9
	61.6	7.6	43.3	34.8	21.4	43.1	4.2
<i>Juniperus occidentalis</i> (bark)	60.1	3.2	49.7	48.1	25.3	31.1	7.1
	–	–	–	–	–	–	–

–, species or component not sampled.

stage of phenology (Table 3) when the goats consumed nearly equal proportions of foliage and bark. Crude protein content of juniper foliage was 7.6% at this time, and an earlier sampling of juniper bark revealed an even lower CP content of 3.2% (Table 4). Juniper bark had the lowest IVDMD value at 31.1%.

Pronounced selective feeding behavior of goats has been observed for legumes (Giger et al., 1987; Morand-Fehr et al., 1987) and also pasture hay and straw (Masson et al., 1989). Degree of selection by goats depends on the plant species offered and stage of growth of the forage. Most often the nutritive value of the forage actually ingested is substantially higher than that of the forage on offer (Morand-Fehr et al., 1980). In natural pastures, there is a high degree of variability in the chemical composition of the ration taken by goats throughout the year (Fedele et al., 1988; Pizzillo et al., 1988). Their selective grazing behavior, however, buffers to a large degree the annual variation in the chemical composition of the pastures.

Goat production in rangeland settings is increasing and resulting in well adapted grazing management techniques. Goats have been utilized in management schemes to keep woody plants at tolerable levels by controlling regrowth subsequent to other brush control measures in the southern California chaparral (Green et al., 1978; Sidahmed et al., 1981). Goats also assist in woody plant control in Tanzania's Massailand (Martin and Huss, 1981), in the northern and central rangelands of Mexico (L.C. Fierro, F. Gomez and M.H. Gonzalez, personal communication, 1980), in reducing Gambel oak (*Quercus gambelii*) sprouts in Colorado (Davis et al., 1975), for woody plant control in western Texas where Spanish goats are used in conjunction with prescribed burning during the growing season (Ueckert, 1980), and in biological control efforts in Australia and New Zealand (Holmst, 1980; Gray, 1984; Radcliffe, 1985). All of these systems, however, require an understanding of feeding behavior, grazing effects on the subsequent growth dynamics and forage values of plants, and the dynamic relationship between the animal and the vegetation. The primary advantage of goats over other ruminants lies in their multipurpose utility (meat, fiber, and milk) and dietary adaptability (Skiles, 1984).

In this study, dietary preference shifted with changes in plant phenology. With our high levels of available forage (530–572 kg ha⁻¹), diverse array of species ($n = 25$), relatively high forage quality and low stocking rates, we observed little potential for control of established Wyoming big sagebrush or western juniper. Our animals secured approximately 99% and 91% of their total bites from the herbaceous component of the vegetation during active growth and dormant phases of plant growth, respectively. Richman et al. (1994) attempted to precondition Angora goats to Wyoming big sagebrush to stimulate later consumption of the shrub in a similar environment. Their sampling occurred principally in 1991, which received roughly 90% of annual precipitation, and about 81% of the selected diet was derived from grasses. Prior exposure to sagebrush had little effect, with no substantial differences between exposed and naive animals, and their goats consumed on average about 1.23 g h⁻¹ of sagebrush in a 15 month trial. About 80% of the sagebrush consumed by their animals consisted of dead material (Richman et al., 1994), which would have little if any impact on subsequent shrub vigor.

In biological control efforts we ideally wish to affect target plants without undue impact on the more desirable crop or forage (Vallentine, 1974). This is especially desirable on federally owned rangelands in the United States where managing agencies typically allow removal of only 40–50% of the available herbage, and there is a strong desire to perpetuate the native flora. Some exemptions to this may be allowed where weed infestations are severe. Whether goats would graze sagebrush or juniper on severely depleted rangelands where selective opportunity is more restricted or on pastures where forage availability is decreased by snow cover is still to be investigated. Likewise, we should research the potential of goats for control of sagebrush and juniper in their early stages establishment. In general, long term trials are still necessary to develop land management strategies with goats in the arid regions of Pacific Northwest. There is need to determine plant responses to various intensities and seasons of grazing, particularly within the little investigated forb component that is not heavily utilized by cattle or horses (McInnis and Vavra, 1987). In closing, however, we find that Spanish goats have little potential

for control of established *Artemisia* and juniper in good condition sagebrush steppe pastures supporting a diverse array of readily available and nutritious forages. Further study is needed, however, to examine their possibilities under an array of conditions where selective grazing opportunities are more restricted or in areas where sagebrush and juniper are in the initial stages of establishment.

Acknowledgements

The Eastern Oregon Agricultural Research Center, including the Burns and Union stations, is jointly operated by the Oregon Agricultural Experiment Station of Oregon State University and the USDA-ARS. This project comprised part of B. Fajemisin's post-doctoral research, concerning the potential of goats as brush control agents. Special thanks are extended to the Center for providing this opportunity. This is Technical Paper 10573, Oregon Agricultural Experiment Station.

References

- Altmann, J., 1974. Observational study of behavior: sampling methods. *Behavior*, 49: 227–265.
- Association of Official Analytical Chemists, 1980. *Official Methods of Analysis*. AOAC, Washington, DC, 1018 pp.
- Austin, D.D., Stevens, R., Jorgensen, K.R. and Urness, P.J., 1994. Preferences of mule deer for 16 grasses found on intermountain winter ranges. *J. Range Manage.*, 47: 308–311.
- Bastian, C.T., Jacobs, J.J. and Smith, M.A., 1995. How much sagebrush is too much: An economic threshold analysis. *J. Range Manage.*, 48: 73–80.
- Britton, C.M. and Sneva, F.A., 1979. Effects of tebuthiuron on a western juniper community. Research in beef cattle nutrition and management. Spec. Rep. 534, Agricultural Experiment Station, Oregon State University, Corvallis.
- Cook, C.W. and Stoddart, L.A., 1960. Physiological responses of big sagebrush to different types of herbage removal. *J. Range Manage.*, 13: 14–16.
- Davis, G.G., Bartel, L.E. and Cook, C.W., 1975. Control of Gambel oak sprouts by goats. *J. Range Manage.*, 28: 216–218.
- Fedele, V., Zarriello, G. and di Trana, A., 1988. Preliminary observations on pasture grazing of local goats. *Agric. Ricerca*, 81: 63–65.
- Giger, S., Sauvant, D., Hervieu, J. and Dorleans, M., 1987. Nutritive value of lucerne hay for goats. *Ann. Zootech.*, 36: 139–152.
- Goering, H.K. and van Soest, P.J., 1970. *Forage Fiber Analysis (Apparatus, Reagents, Procedures and some Applications)*. Agricultural Handbook 379. ARS, USDA, Washington DC, 20 pp.
- Gray, T., 1984. Goats curb weed problem. *N.Z. Farmer*, 105: 13–14.
- Green, L.R., Hughes, C.L. and Graves, W.L., 1978. Goat control of brush regrowth on southern California chaparral fuel breaks. In: *Proc. 1st Int. Rangeland Congress*, 14–18 August 1978, Denver, CO. Society for Range Management, Denver, CO, pp. 451–455.
- Hitchcock, C.L. and Cronquist, A., 1973. *Flora of the Pacific Northwest*. University of Washington Press, Seattle, 730 pp.
- Holechek, J.L., Vavra, M. and Pieper, R.D., 1984. Methods of determining the botanical composition, similarity and overlap of range herbivore diets. In: *Developing Strategies for Rangeland Management*. National Research Council (NRC) National Academy of Sciences, Westview Press, Boulder, CO, pp. 425–471.
- Holmst, P.J., 1980. The use of goats in grazing systems and their place in weed control. *Proc. Aust. Soc. Anim. Prod.*, 13: 188–191.
- Jones, T.A., Ralphs, M.A. and Nielson, D.C., 1994. Cattle preference for 4 wheatgrass taxa. *J. Range Manage.*, 47: 119–122.
- Leclerc, B., 1984. Utilization of Corsican chaparral by goats and sheep. I—Goat's diet. *Acta Oecol. Ecol. Applic.*, 5: 383–406.
- Malechek, J.C. and Provenza, F.D., 1981. Feeding behavior and nutrition of goats on rangelands. In: P. Morand-Fehr, A. Bourbouze and M. de Simiane (Editors), *Nutrition and Systems of Goat Feeding*, Tours, France, 12–15 May, INRA-ITOVIC, Paris, Vol. 1, pp. 411–428.
- Martin, J.A. and Huss, D.L., 1981. Goats much maligned but necessary. *Rangelands*, 3: 199–201.
- Masson, C., Kirilov, D., Faurie, F. and Tisserand, J.L., 1989. Feeding behavior and rumination in sheep and goats fed with hydroxide treated or untreated barley straw. *Ann. Zootech.*, 38: 73–82.
- McInnis, M.L. and Vavra, M., 1987. Dietary relationships among feral horses, cattle and pronghorn in southeastern Oregon. *J. Range Manage.*, 40: 60–66.
- Miller, R.F., Svejcar, T.J. and West, N.E., 1994. Implication of livestock grazing in the intermountain sagebrush region: plant composition. In: M. Vavra, W.A. Laycock and R.D. Piper (Editors), *Ecological Implications of Livestock Herbivory in the West*. Society for Range Management, Denver, CO, pp. 101–146.
- Morand-Fehr, P., 1988. Capacity of goat adaptation in arid environments. In: *La Chevre. Ethnozootechnie*. Societe d'Ethnozootechnie, Paris, Vol. 41, pp. 63–85.
- Morand-Fehr, P., Hervieu, J. and Sauvant, D., 1980. Contribution to the description of feeding pattern. *Reprod. Nutr. Develop.*, 20: 1641–1644.
- Morand-Fehr, P., Giger, S., Sauvant, D., Broqua, B. and de Simiane, M., 1987. Dry forage utilization for goats. In: C. Demarquilly (Editor), *Les Fourrages Secs 1: Recolte, Traitement, Utilization*. INRA, Paris, pp. 391–422.
- Narjisse, H. and El Bare, B., 1986. Seasonal changes in the

- dietary botanical composition of sheep and goats grazing in an oak forest. In: P.J. Joss, P.W. Lynch and O.B. Williams (Editors), *Rangelands: A Resource Under Siege*. Australian Academy of Science, Canberra, pp. 369–371.
- Pizzillo, M., di Trana, A. and Fedele, V., 1988. Grazing feeding behavior of Maltese breed and local goats. *Agric. Ricerca*, 81: 69–78.
- Pringle, W.L., 1960. The effect of a leaf feeding beetle on big sagebrush in British Columbia. *J. Range Manage.*, 13: 139–142.
- Radcliffe, J.E., 1985. Grazing management of goats and sheep for gorse control. *N.Z. J. Exp. Agric.*, 13: 181–190.
- Richman, L.M., 1993. Diet selection by conditioned and unconditioned goats in the sagebrush steppe of eastern Oregon. M.S. Thesis, Oregon State University, Corvallis, 98 pp.
- Richman, L.M., Johnson, D.E. and Angell, R.F., 1994. Evaluation of a positive conditioning technique for influencing big sagebrush (*Artemisia tridentata* subsp. *wyomingensis*) consumption by goats. *Appl. Anim. Behav. Sci.*, 40: 229–240.
- Rubino, R., Pizzillo, M. and Scardella, P., 1988. Comparison between different grazing systems in an intensive breeding model of Maltese goats. *Agric. Ricerca*, 81: 79–86.
- Schwartz, C.C., Nagy, J.G. and Rice, R.W., 1977. Pronghorn dietary quality relative to forage availability and other ruminants in Colorado. *J. Wildl. Manage.*, 41: 161–168.
- Sidahmed, A.E., Morris, J.G. and Radosevitch, S.R., 1981. Summer diet of Spanish goats grazing chaparral. *J. Range Manage.*, 34: 33–35.
- Skiles, J.W., 1984. A review of animal preference. In: *Developing Strategies for Rangeland Management*. National Research Council/National Academy of Science, Westview Press, Boulder, CO, pp. 153–213.
- Sneva, F.A., 1982. Relation of precipitation and temperature with yield of herbaceous plants in eastern Oregon. *Int. J. Biometeorol.*, 26: 263–276.
- Tilley, J.M.A. and Terry, R.A., 1963. A two-stage technique for the in-vitro digestion of forage crops. *J. Br. Grassl. Soc.*, 18: 104.
- Ueckert, D.N., 1980. Manipulating range vegetation with prescribed burning. *Proc. Symp. on Prescribed Range Burning in the Edwards Plateau of Texas, 23 October 1980, Junction, TX*. Texas Agricultural Extension Service, College Station, TX, pp. 27–44.
- Vallentine, J.F., 1974. *Range Development and Improvements*. Brigham Young University Press, Provo, 516 pp.
- Wright, H.A., 1970. Response of big sagebrush and three-tip sagebrush to season of clipping. *J. Range Manage.*, 23: 20–21.